



## King's Research Portal

[Link to publication record in King's Research Portal](#)

### *Citation for published version (APA):*

Ridsdale, L. L., & O'Flynn, N. (2002). Headache. In M. Marshall, S. Campbell, J. Hacker, & M. Roland (Eds.), *Quality indicators for general practice* (pp. 112). Royal Society of Medicine Press Ltd.

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

Norma O'Flynn and Leone Ridsdale

[illegible]

### Review of evidence relating to quality indicators

### Diagnosis

Headaches are largely diagnosed as clinical syndromes. Primary headaches are those for which there is no structural or metabolic cause. The objective of the diagnostic process is to rule out secondary causes, particularly serious causes of headache, and then to accurately diagnose primary headache to ensure appropriate treatment.

**History**  
The most common primary headaches are tension headache and migraine. Secondary headache is most commonly associated with excess alcohol consumption, medication misuse, fever and disorders of the nose and sinuses.<sup>1</sup> Headache is diagnosed by focusing on the exact definition of the prominent symptoms, in particular the temporal pattern, the description of pain and associated signs and symptoms, as well as related phenomena. Primary headaches can coexist: a patient may have tension headache and migraine.

headache itself need to be assessed, such as the location (migraine is usually unilateral but 30–40% can be bilateral), the quality (e.g. pulsating, pressure), the clinical features of classical migraine by movement, Prodromal features or aura are sensitivity to light and sound predominantly occur with migraine. Nausea or vomiting during an attack, and headache history include family history, precipitating factors. Other features of factors and a drug history.

Aspects of the history that may indicate underlying pathology are new-onset headache, particularly in middle age or later, worst-ever headache, changes in frequency or severity of usual features of previous long-standing headache, changes in presence of systemic features such as fever, anorexia, weight loss, and the claudication.<sup>9</sup>

There is very little literature on the sensitivity and specificity of aspects of the history. In a Danish population, a questionnaire validated against a telephone interview found that the two questions 'Have you ever had migraine?' and 'Have you ever had visual disturbances lasting 5–60 min followed by headache?' identified 93% of migraineurs with aura and 75% of migraineurs without aura.<sup>1</sup> The findings are likely to differ in populations with less knowledge of migraine. Jaw claudication is strongly suggestive of giant cell arteritis; the odds of a positive biopsy are 9 times greater if jaw claudication is present.<sup>2</sup> Individual items of the clinical history have low accuracy for the diagnosis of meningitis in adults. Results from a review of studies with meningitis indicated sensitivity of 50% for headache (CI 32–68%) and sensitivity of 30% for nausea/vomiting (CI 22–38%).<sup>3,2</sup>

There is lack of agreement about the essential elements of the physical examination. If the main symptom is headache, and the history has not revealed any focal symptoms, complete physical and neurological examination is not usually helpful.<sup>9</sup> Examinations suggested by neurologists include blood pressure and extracranial structures such as sinuses, scalp arteries, cervical paraspinal muscles and temporomandibular joints (TMJs). Suggested neurological examination includes optic fundi, cranial nerves (in particular the Vth cranial nerve, including the corneal reflex); muscle power, gait, reflexes and plantar responses. The prime objective of the examination is to support the diagnostic hypothesis suggested by the history and to exclude causes of headache that require further investigation.

There is a lack of evidence as to the precise value of most elements of the examination, and no evidence as to their predictive value in a general practice population. A questionnaire designed in an American headache centre to screen for temporomandibular disorders found high sensitivity and specificity (92% and 91% respectively) for pain with maximum jaw opening using passive stretch and reciprocal jaw clicking or pain on palpitation over TMJ, to distinguish TMJ derangement from headache patients.<sup>19</sup> One case control study in a headache centre found that papilloedema was an uncommon sign in acute elevation of intracranial pressure.<sup>20</sup>

[illegible]

Laboratory investigations such as Hb, CRP and ESR are suggested when the diagnosis of temporal arteritis is suspected. However, evidence from the literature is conflicting. One retrospective study of patients referred for biopsy found no significant difference in either parameter in biopsy-positive or biopsy-negative cases.<sup>25</sup> However, a study using controls as well as the group referred for biopsy found a sensitivity of 100% for CRP, 92% for ESR and the combination of ESR and CRP was highly specific (97%) for temporal arteritis.<sup>11</sup>

**Referral (indicator 1)**  
The Referral Guidelines for Suspected Brain Tumour suggest that a referral should be made and patient seen within 2 weeks for:

- ▶ subacute progressive focal neurological deficit (e.g. weakness, sensory loss, dysphasia, ataxia)
- ▶ new-onset seizures characterised by one or more of
  - focal seizures
  - prolonged (greater than 1 hour) post-ictal deficit
  - status epilepticus
  - associated interictal focal deficit
- ▶ patients with headache, vomiting and papilloedema,
- ▶ cranial nerve palsy (e.g. diplopia, visual failure including optician-defined visual field loss, unilateral sensorineural deafness).

Urgent referral should be considered for

- Urgent referral should be considered for
- ▶ headache patients with non-migrainous headaches of recent onset, present for at least 1 month, when accompanied by features suggestive of raised intracranial pressure (e.g. woken by headache, vomiting, drowsiness)

This section deals with the evidence for the treatment of primary headache. We have confined our comments to therapeutic agents likely to be used in a UK primary care context. Experts comment on the importance of eliminating triggers, that stress can precipitate any type of headache and that non-drug treatment such as relaxation should be considered.

In the case of migraine, treatment options vary as to the severity of the acute attack. Limmoth et al.<sup>21</sup> reviewed the use of aspirin (acetylsalicylic acid or ASA) in acute treatment and prophylactic treatment of migraine. In the acute phase its effectiveness is enhanced by an effervescent form and combination with antiemetics (of note multicentre trial, water-soluble aspirin with metoclopramide was as effective as 50 mg of oral sumatriptan in reducing headache intensity and autonomic features such as nausea, and it had fewer adverse effects.<sup>22</sup> Ibuprofen, naproxen and diclofenac sodium have been shown to be effective treatments for an acute attack.<sup>23-25</sup> 50 mg of oral diclofenac K provided more rapid relief than oral sumatriptan in a double-blind randomised crossover trial.<sup>26</sup> (The potassium salt is more rapidly absorbed than the sodium.)

Triptan use is recommended for moderate or severe migraine. A systematic review found that subcutaneous sumatriptan was more efficacious than the oral form and had a quicker onset of action, but caused more side-effects.<sup>29</sup> The investigator stated that the majority of side-effects were minor and may be tolerable to patients. A prospective study of the tolerability of subcutaneous sumatriptan in acute migraine, which followed 12 239 patients over 12 months, found no major adverse effects when sumatriptan was used according to the precautions and warnings on the label.<sup>30</sup>

There is evidence from randomised controlled trials for the effectiveness of intramuscular chlorpromazine, metoclopramide and prochlorperazine for an acute attack.<sup>31,32</sup>

Overuse of most acute attack therapies can aggravate headache frequency, and particular care needs to be taken with codeine-containing compound analgesics.

Prophylactic treatment is suggested if attacks occur frequently, or if attacks severely incapacitate the patient (indicator 2). Traditionally, experts have suggested that prophylaxis should be considered if a patient suffers 2 attacks or more each month, but more effective remedies for the acute attack may vary this. The efficacy of commonly used prophylactic drugs is low.<sup>13</sup>

There is clinical trial evidence for the effectiveness of the following drugs in prophylaxis (only propranolol and pizotifen are licensed for migraine prophylaxis in the UK): beta-blockers (atenolol, metoprolol, naldolol, propranolol);<sup>31,32</sup> calcium channel blockers (verapamil, flunarizine);<sup>33,34</sup> serotonin receptor antagonists (pizotifen, methysergide);<sup>35,36</sup> tricyclics (amitriptyline);<sup>37</sup> and 'antiepileptics' (sodium valproate, valproic acid)<sup>32,43</sup> (indicator 3).

There is evidence for the effectiveness of naproxen but the expert recommendation is to confine its use to the prophylaxis of perimenstrual migraine as intermittent use results in fewer gastrointestinal side-effects.<sup>46,45</sup> Expert opinion is that failure of one

beta-blocker does not necessarily predict the response to another. Treatment should start with a low dose and titrate upwards; once pain is controlled, the dose may be tapered down.

In the treatment of headache other than migraine, aspirin and ibuprofen have been shown to be effective in the treatment of episodic tension-type headache.<sup>38</sup> Dosage may be important: a double-blind placebo-controlled comparison of ketoprofen 25 mg and paracetamol 1000 mg found that both were more effective than placebo, with no significant difference between the active agents;<sup>41</sup> and a comparison of ibuprofen 400 mg with paracetamol 1000 mg found ibuprofen to be the more effective.<sup>42</sup> Tricyclics have been found to be effective in reducing the frequency and severity of chronic tension-type headaches.<sup>43</sup>

There is randomised controlled trial evidence for the effect of spinal manipulation in the case of cervicogenic headache.<sup>39</sup> However, spinal manipulation does not have a positive effect on episodic tension-type headache.<sup>41</sup>

Certain prophylactic agents should not be prescribed in specific at-risk groups, for example sumatriptan for patients with angina and beta-blockers for asthmatics (indicator 4).

#### Follow-up

Experts recommend the use of headache diaries in the assessment and treatment of headache.<sup>3,45</sup> Prophylactic treatment also requires regular monitoring.

#### Headache in children and adolescents

The diagnosis and treatment of headache in children is broadly similar to that in adults. The criteria for the diagnosis of migraine in children is different, in part relating to aspects of the headache itself, but also noting the difficulty that children may have in articulating characteristics such as the intensity and the nature of pain.<sup>51</sup> Mortimer et al.<sup>52</sup> interviewed children registered at one general practice. Headache prevalence increased from the age of 3 years up to age 11 in boys and girls, with a higher prevalence in 3 to 5-year-old boys than 3 to 5-year-old girls. The overall prevalence of headache was 38.8%, and of migraine was 3.7%. There is a suggestion that tension-type headache in children is associated with psychosocial problems, but the evidence from the literature is conflicting.<sup>53</sup> The history and examination should be supplemented by a developmental assessment. Simple analgesia such as paracetamol is suggested, used early and in adequate dosage (Committee on Safety of Medicines advice is to avoid the use of aspirin in children under 12 years, except for specific indications, due to the risk of Reye's syndrome). Metoclopramide can cause severe extrapyramidal side-effects in children. Zolmitriptan is licensed for use from the age of 12. Pizotifen is used for prophylaxis.

#### Recommended quality indicators for headache

##### Diagnosis/referral

- 1 Patients should be referred urgently for specialist care and investigation if the presenting headache is accompanied by:
  - a. suspected raised intracranial pressure
  - b. new-onset seizure
  - c. focal neurological signs
  - d. papilloedema

##### Treatment

- 2 Prophylaxis treatment should be offered in patients with severe and disabling migraine
- 3 The following agents should be prescribed as first line for prophylaxis of migraine unless contraindicated:
  - a. beta-blocker
  - b. tricyclic antidepressant
  - c. pizotifen
- 4 a. Sumatriptan should not be prescribed for migraine in patients with angina  
b. Beta-blockers should not be prescribed for migraine in patients with asthma

#### Overview of data sources used in this review

Articles on headache were identified by a search of the 1991–5 and 1996–9 Medline databases and the Cochrane library. Keywords in the search were 'headache' (exploded and used as textword), 'family practice', 'epidemiology', 'sensitivity and specificity', and 'treatment'. We searched particularly for literature with a community/general practice focus. However, the majority of evidence is from specialist centres and uses the classification of headache of The International Headache Society (IHS).<sup>56</sup> This establishes diagnostic criteria for headache using a hierarchical scale and has been used since 1988 for epidemiological and clinical research.<sup>56</sup> Guidelines for the diagnosis and management of headache and migraine have been produced by a number of groups and these were also reviewed.<sup>57,58</sup> Migraine management guidelines have been produced by the British Association for the Study of Headache<sup>59</sup> and a group endorsed by Migraine in Primary Care Advisors.<sup>60</sup> Referral Guidelines for Suspected Brain Tumours have been prepared as part of the initiative to allow patients with suspected cancer to be seen by a specialist within 2 weeks.

There is an absence of evidence relating to the management of headache in primary care practice and many aspects of diagnosis and treatment are based on expert opinion.

#### Further reading

Goadsby Peter J, Olesen Jes. Fortnightly review: diagnosis and management of migraine. *British Medical Journal* 1996; 312: 1279–1283



Pryse-Phillips WE, Dodick DW, Edmeads JC et al. Guidelines for the diagnosis and management of migraine in clinical practice. *Canadian Medical Association Journal* 1997; 156: 1273-1287  
The British Association for the Study of Headache (BASH) has a website at [www.bash.org.uk](http://www.bash.org.uk)

## References

- Rasmussen BK, Jensen R, Schroll M, Olesen J. Epidemiology of headache in a general population - a prevalence study. *Journal of Clinical Epidemiology* 1991; 44(11): 1147-1157
- Lipton RB, Stewart WF. The epidemiology of migraine. *European Neurology* 1994; 34(suppl 2): 6-11
- Rasmussen BK. Epidemiology of headache. *Cephalalgia* 1995; 15: 45-68
- Jensen R. Pathophysiological mechanisms of tension type headache: a review of epidemiological and experimental studies. *Cephalalgia* 1999; 19: 602-621
- Scher AI, Stewart WF, Liberman J, Lipton RB. Prevalence of frequent headache in a population sample. *Headache* 1998; 38: 497-506
- Castillo J, Munoz P, Gutera V, Pascual J. Epidemiology of chronic daily headache in the general population. *Headache* 1999; 39: 190-196
- Royal College of General Practitioners, Office of Populations Censuses and Surveys, Department of Health. Morbidity statistics in general practice fourth national study 1991-1992. London: HMSO, 1995
- Cockrell OC, Goodridge DMG, Brodie D, Sander JWAS, Shorvon SD. Neurological disease in a defined population: the results of a pilot study in two general practices. *Neuroepidemiology* 1996; 15: 73-82
- Goadsby Peter J, Olesen J. Fortnightly review: diagnosis and management of migraine. *British Medical Journal* 1996; 312: 1279-1283
- Gervil M, Ulrich V, Olesen J, Russell MB. Screening for migraine in the general population: validation of a simple questionnaire. *Cephalalgia* 1998; 18: 342-348
- Hayreh SS, Podhajsky PA, Raman R, Zimmerman B. Giant cell arteritis: validity and reliability of various diagnostic tests. *American Journal of Ophthalmology* 1997; 123(3): 285-296
- Atia J, Hatala R, Cook DJ, Wong JG. Does this adult patient have acute meningitis? *Journal of the American Medical Association* 1999; 282: 175-181
- Breslau N, Schultz LR, Stewart WF, Lipton RB, Lucia VC, Welch KMA. Headache and major depression. Is the association specific to migraine? *Neurology* 2000; 54: 308-313
- Holroyd KA, Stensland M, Lipchik GL, Hill KR, O'Donnell FS, Cordingley G. Psychosocial correlates and impact of chronic tension-type headaches. *Headache* 2000; 40: 3-16
- Mitskostas DD, Thomas AM. Comorbidity of headache and depressive disorders. *Cephalalgia* 1999; 19: 211-217
- Wang SJ, Liu HC, Fuh JL, Liu CY, Wang PN, Lu SR. Comorbidity of headache and depression in the elderly. *Pain* 1999; 82(3): 239-243
- Ham LP, Andarsk F, Packard RC, Burdick CM. Psychopathology in individuals with post-traumatic headache and other pain types. *Cephalalgia* 1994; 14: 118-126
- Marks DR, Papoport AM. Practical evaluation and diagnosis of headache. *Seminars in Neurology* 1997; 17(4): 307-312
- Schiffman E, Haley D, Baker C, Lindgren B. Diagnostic criteria for screening headache patients for temporomandibular disorders. *Headache* 1995; 35: 121-124
- Steffen H, Ertel B, Aschoff A, Kolling GH, Volcker HE. The diagnostic value of optic disc evaluation in acute elevated intracranial pressure. *Ophthalmology* 1996; 103(8): 1229-1232
- Frisberg B. Neuroimaging in presumed primary headache disorders. *Seminars in Neurology* 1997; 17(4): 373-382
- Chmielewski WL, McKnight KM, Agudelo CA, Wise CM. Presenting features and outcomes in patients undergoing temporal artery biopsy. *Archives of Internal Medicine* 1992; 152: 1690-1695
- Limmroth V, Katsarva Z, Diener H-C. Acetylsalicylic acid in the treatment of headache. *Cephalalgia* 1999; 19: 545-551
- Tfelt-Hansen P, Henry P, Mulder LJ, Schneidewert RG, Schoonen J, Chazot G. The effectiveness of combined oral lysine acetylsalicylate and metoprolol compared with oral sumatriptan for migraine. *Lancet* 1995; 346: 925-926
- Treves TA, Streiffert M, Korczyn AD. Naproxen sodium versus ergotamine tartrate in the treatment of acute migraine attacks. *Headache* 1992; 32(6): 280-282
- Karachalios GN, Fotiadou A, Nickolaou G, Karabetsos A, Kefalogiorgou K. Treatment of acute migraine attack with diclofenac sodium: a double-blind study. *Headache* 1992; 32: 88-100
- Kloster R, Nestvold K, Vilming ST. A double-blind study of ibuprofen versus placebo in the treatment of acute migraine attacks. *Cephalalgia* 1992; 12: 169-171
- Diclofenac-K-Sumatriptan Migraine Study Group. Acute treatment of migraine: efficacy and safety of a non-steroidal anti-inflammatory drug, diclofenac-potassium in comparison to oral sumatriptan and placebo. *Cephalalgia* 1999; 19: 232-240
- Tfelt-Hansen P. Efficacy and adverse effects of subcutaneous, oral and intranasal sumatriptan used for migraine treatment: a systematic review based on number needed to treat. *Cephalalgia* 1998; 18: 532-538
- O'Quinn S, Davis RL, Guterman DL, Pait GD, Fox AW. Prospective large-scale study of the tolerability of subcutaneous sumatriptan injection for the acute treatment of migraine. *Cephalalgia* 1999; 19: 223-231
- Lane PL, McLellan BA, Baggoley CJ. Comparative efficacy of chlorpromazine and meperidine with dimenhydrinate in migraine headache. *Annals of Emergency Medicine* 1989; 18(4): 360-365
- Coppola M, Yealy DM, Laibold RA. Randomised, placebo controlled evaluation of prochlorperazine versus metoprolol for emergency department treatment of migraine headaches. *Annals of Emergency Medicine* 1995; 26: 541-546
- Ramadan NM, Schultz LL, Gilkey SJ. Migraine prophylactic drugs: proof of efficacy, utilization and cost. *Cephalalgia* 1997; 17(2): 73-80
- Johannsson V, Nilsson LR, Widelius T et al. Atenolol in migraine prophylaxis: a double blind cross over multicentre study. *Headache* 1987; 27: 372-4
- Gerber WD, Diener H C, Scholz E, Niederberger U. Responders and non-responders to metoprolol, propranolol and nifedipine treatment in migraine prophylaxis: a dose range study based on time series analysis. *Cephalalgia* 1991; 11: 37-45
- Ryan SR, Ryan JR, Sudilovsky A, Nadolol: its use in the prophylactic treatment of migraine. *Headache* 1983; 23: 26-31
- Markley HG. Verapamil and migraine prophylaxis: mechanisms and efficacy. *American Journal of Medicine* 1991; 90(suppl 5A): 48S-53S
- Soelberg Sorensen P, Larsen BH, Rasmussen MJ, King E, Iversen H. Flunarizine versus metoprolol in migraine prophylaxis: a double blind, randomised parallel group study of efficacy and tolerability. *Headache* 1991; 31: 650-657
- Drummond PD. Effectiveness of methysergide in relation to clinical features of migraine. *Headache* 1985; 5: 145-146
- Capitolo R, Rose FC. Single dose pizotifen, 1.5 mg nocte: a new approach in the prophylaxis of migraine. *Headache* 1982; 22: 272-275
- Bank J. A comparative study of amitriptyline and fluvoxamine in migraine prophylaxis. *Headache* 1994; 34: 476-479
- Herring R, Kuritzky RS. Sodium valproate in the prophylactic treatment of migraine: a double blind study versus placebo. *Cephalalgia* 1992; 12: 81-84
- Jensen R, Brinck T, Olesen J. Prophylactic effect of sodium valproate in migraine without aura: a double-blind, placebo- and dose-controlled study. *Cephalalgia* 1993; 13(13): 193
- Welch KMA, Ellis DJ, Keenan PA. Successful migraine prophylaxis with naproxen sodium. *Neurology* 1985; 35: 1304-1310
- Szekely B, Merryman S, Croft H, Post G. Prophylactic effects of naproxen sodium on perimenstrual headache: a double blind placebo controlled study. *Cephalalgia* 1989; 9 (suppl 10): 452-453
- Nebe J, Heier M, Diener HC. Low dose ibuprofen in self medication of mild to moderate headache: a comparison with acetylsalicylic acid and placebo. *Cephalalgia* 1995; 15: 531-535
- Steiner TJ, Lange R. Ketoprofen (25 mg) in the symptomatic treatment of episodic tension type headache: double-blind placebo-controlled comparison with acetaminophen (1000 g). *Cephalalgia* 1998; 18(1): 38-43
- Schäpflin BP, Furey SA, Thoden WR. Non-prescription ibuprofen and acetaminophen in the treatment of tension type headache. *Journal of Clinical Pharmacology* 1996; 36 (12): 1120-1125
- Bendtsen L, Jensen R, Olesen J. A non-selective (amitriptyline), but not a selective (citalopram), serotonin reuptake inhibitor is effective in the prophylactic treatment of chronic tension-type headache. *Journal of Neurology, Neurosurgery and Psychiatry* 1996; 61(3): 285-290
- Nilsson NA, Christensen HW, Hartvigsen J. The effect of spinal manipulation in the treatment of cervicogenic headache. *Journal of Manipulative and Physiological Therapeutics* 1997; 20: 326-330
- Bove G, Nilsson N (1998) Spinal manipulation in the treatment of episodic tension-type headache: a randomised controlled trial. *Journal of the American Medical Association* 1998; 280: 1576-1579

## Quality indicators for general practice

- Russell MB, Rasmussen BK, Brennum J, Jensen HK, Jensen RA, Olesen J. Presentation of a new instrument: the diagnostic headache diary. *Cephalalgia* 1992; 12: 369-374
- Lipton RB. Diagnosis and epidemiology of pediatric migraine. *Current Opinion in Neurology* 1997; 10: 231-235
- Mortimer J, Kay J, Jaron A. Epidemiology of headache and childhood migraine in an urban general practice using ad hoc, Vahquist and IHS criteria. *Developmental Medicine and Child Neurology* 1992; 34: 1095-1101
- Karwautz A, Wober C, Lang T et al. Psychosocial factors in children and adolescents with migraine and tension type headache: a controlled study and review of the literature. *Cephalalgia* 1999; 19: 32-43
- Headache Classification Committee of the International Headache Society. Classification and diagnostic criteria for headache disorders, cranial neuralgias and facial pain. *Cephalalgia* 1988; 8 (suppl 7)
- Pryse-Phillips WE, Dodick DW, Edmeads JC et al. Guidelines for the diagnosis and management of migraine in clinical practice. *Canadian Medical Association Journal* 1997; 156: 1273-1287
- Danish Neurological Society and Danish Headache Society. Guidelines for the management of headache. *Cephalalgia* 1998; 18: 9-22
- British Association for the Study of Headache. Guidelines for all doctors in the diagnosis and management of migraine. London: Princess Margaret Migraine Clinic, 1999
- Migraine in Primary Care Advisors. Migraine management guidelines. Synergy Medical Education, 1997

## ► 16

## Upper respiratory tract infections

Paul Little

## Importance

Acute upper respiratory tract infections (URTIs) are the commonest reason for patients to seek medical advice<sup>1</sup> and also the commonest reason for antibiotics to be prescribed. The current major concern is that the inappropriate use of antibiotics for usually self-limiting conditions will foster the development of antibiotic resistance and lead to serious infections becoming untreatable.<sup>2-4</sup> For this reason it is currently a national priority not to encourage the use of antibiotics unless there is very good evidence of their efficacy.

## Pharyngitis/tonsillitis: diagnosis

Pharyngitis is caused by both bacterial and viral organisms.<sup>5</sup> Antibiotics could be targeted to those who have positive throat swabs for group A Streptococcus, a positive rapid Strep. test, or clinical characteristics associated with a positive throat swab.<sup>6-11</sup> Alternatively throat swabs could be used in selected populations.<sup>12</sup> However, throat swabs increase costs significantly, potentially medicalise self-limiting illness<sup>13</sup> and rarely modify clinical decisions.<sup>14</sup> In addition, antibiotic use and overall accuracy of decision making may be unchanged.<sup>14</sup> The throat swab is neither particularly sensitive nor specific when compared to a rise in antistreptolysin-O titres (ASOT). This is the case in both the general population<sup>15</sup> and in selected general practice populations where clinical selection has occurred.<sup>16</sup> Clinical scores or decision rules based on the throat swab<sup>17-19</sup> have the same limitations of validity as the throat swab, although they may crudely identify patients at risk of complications (see below).

## Pharyngitis/tonsillitis: treatment

### Antibiotics

A systematic review<sup>20</sup> indicates that antibiotics reduce symptom duration by a few hours to half a day. For patients who are not systemically unwell, either not prescribing or using a delayed prescribing approach (waiting for several days before using the prescription) is acceptable, changes attitudes to antibiotics, modifies attendance behaviour and does not delay symptom resolution appreciably.<sup>18,19</sup> Delaying the prescription probably results in 20% fewer recurrences compared to the immediate prescriptions of antibiotics, presumably because antibiotics modify local or systemic immune mechanisms.<sup>20-22</sup>